

Chemical composition, antiradical and phytotoxic activity of the essential oil from *Peucedanum ostruthium* W.D.J.Koch leaves

Stefania Garzoli¹ , Marcello Iriti^{2, 4*} , Sara Vitalini^{3, 4*} 

¹Department of Drug Chemistry and Technology, Sapienza University, P.le Aldo Moro 5, 00185, Rome, Italy

²Department of Biomedical, Surgical and Dental Sciences, Università degli Studi di Milano, Via G. Celoria 2, 20133 Milan, Italy

³Department of Food, Environmental and Nutritional Sciences, Università degli Studi di Milano, via Mangiagalli 25, 20133, Milan, Italy

⁴National Interuniversity Consortium of Materials Science and Technology, via G. Giusti 9, 50121 Firenze, Italy

Tel.: +39 02 50316766; fax: +39 02 50316781

Abstract

Peucedanum ostruthium W.D.J.Koch, commonly known as masterwort, is a flowering perennial species in the Apiaceae family with known medicinal and aromatic properties. This study was designed to chemically characterize the essential oil (EO) obtained from the leaves and investigate some aspects of its bioactivity. Thirty-two compounds were detected by gas chromatography-mass spectrometry analysis and sesquiterpenoids identified as the dominating group of compounds. The major ones were caryophyllene oxide (20.7%) and spathulenol (17.2%), followed by cubenol (8.7%), δ-cadinene (6.1%) and humulene epoxide II (5.6%). EO was evaluated *in vitro* by ABTS⁺ (2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) and DPPH[·] (2,2-diphenyl-picryl hydrazyl) assays, showing a marked scavenging ability, in particular towards the ABTS⁺ radical cation (2.02 ± 0.00 μM Trolox eq/mL). EO was also screened for phytotoxic activity against mono- and dicotyledonous weeds. It exhibited significant effects by reducing the growth of *Lolium multiflorum* Lam. and *Sinapis alba* L. seedlings up to 90.7% and 76.6%, respectively.

Article Information

Received: 15 November, 2022

Revised: 28 November, 2022

Accepted: 12 December, 2022

Academic Editor

Ayşe Kuruüzüm-Uz

Corresponding Authors

Marcello Iriti

E-mail:

marcello.iriti@unimi.it

Sara Vitalini

E-mail:

sara.vitalini@unimi.it

Keywords

Masterwort, Apiaceae, allelopathy; radical scavenging effects, GC-MS analysis, VOCs

1. Introduction

Peucedanum ostruthium W.D.J.Koch (syn. *Imperatoria ostruthium* L.), commonly referred to as masterwort, is a perennial herbaceous species belonging to the Apiaceae family. Its native range is the mountains of central and southern Europe. In Italy, it grows spontaneously in the central and northern regions between 1000 and 2000 m. a.s.l. *P. ostruthium* is characterized by white flowers collected in umbrellas, an erect stem up to 40 cm high and tripartite leaves with a deeply serrated edge [1, 2]. Flowers, leaves and rhizomes are used in folk medicine for remedies administered both internally and externally to treat cardiovascular, digestive, integumentary, musculo-skeletal, respiratory and urogenital disorders. The

rhizome is also appreciated for its flavoring properties exploited in the preparation of some drinks such as grappas and liqueurs [3-5].

Peucedanum species are known to possess a broad spectrum of pharmacological activities and many of them are due to the presence of coumarins, flavonoids, phenolics and essential oils [6]. The latter were investigated for their antimicrobial, antioxidant and cytotoxic [7-10] as well as pesticidal [11] effects obtaining promising results. The bioactivity of *P. ostruthium* essential oil (EO) has never been studied. Therefore, this work was designed to examine its antiradical and phytotoxic activity, after the characterization of the chemical composition.



2. Materials and methods

2.1 Plant material collection

The leaves of *P. ostruthium* were collected in August 2018 in the Valle di Champorcher (Valle d'Aosta, Italy) at 1955 m a.s.l and air dried, then stored in paper bags until pulverization with a laboratory blender. The species was identified according to Flora d'Italia [2] and the herbarium sample (No. PO-VC-VDA-18) was deposited at the Department of Agricultural and Environmental Sciences of the Milan State University (Italy).

2.2 Seeds

The seeds of *S. alba* were purchased from the company "Padana Sementi" situated in Tombolo (Padova, Italy) while those of *L. multiflorum* were supplied by the organic rice farm "Terre di Lomellina", located in Candia Lomellina (Pavia, Italy). After an appropriate selection, they were surface sterilized with a 1% bleach solution for 10 minutes, then rinsed repeatedly with distilled water until the disinfectant was completely removed.

2.3 Essential oil distillation

EO of the dried leaves from *P. ostruthium* was obtained by 3 h hydrodistillation in a Clevenger-type circulatory apparatus with 25 g of material and a yield of 0.68% w/w. Then, it was collected and combined with anhydrous sodium sulfate to remove water. EO was stored at 4 °C in a sealed vial until use.

2.4 Chemical analysis

To perform the analyses, a Clarus 500 model Perkin Elmer (Waltham, MA, USA) gas chromatograph coupled with a mass spectrometer and equipped with an FID (flame detector ionization) was used. Chromatographic separation was performed with a Varian Factor Four VF-1 capillary column flushed with helium at a flow rate of 1 mL/min. The applied operative conditions followed those reported in some previous works [12,13]. The mass spectra were obtained in the electron impact mode (EI), at 70 eV in scan mode in the range 35-400 m/z. The identification of volatile compounds was performed by matching their mass spectra with those stored in the Wiley 2.2 and Nist 02 mass spectra libraries database and by comparison of their linear retention indices (LRIs),

relative to C8-C25 *n*-alkanes, with those available in the literature. To express the quantity of the components, we used the percentage values calculated in relation to the total area of the chromatogram by normalizing the peak area without the use of an internal standard and any factor correction. The analysis was carried out in triplicate.

2.5 Antiradical assays

DPPH assay

The radical-scavenging capacity of EO against DPPH- was assessed following Iriti et al. [14], with some modifications. Briefly, the DPPH- stock solution (0.35 g/L) was diluted with methanol to an absorbance of 1.00 ± 0.03 at 517 nm. Then, 50 μ L of EO was added to 2.45 mL of this solution. After a reaction time of 30 min in the dark and at room temperature, the absorbance changes were monitored using a Jenway 6310 spectrophotometer (Keison, Chelmsford, Essex, UK). The obtained results are expressed as μ M eq Trolox mL⁻¹ EO. A DPPH- solution without EO was used as a control. The assay was performed in triplicate.

ABTS assay

The test was carried out following Iriti et al. [14]. The ABTS⁺ radical cation was produced by reacting a solution of ABTS 7 mM with potassium persulfate 2.45 mM and keeping the mixture in the dark at room temperature for at least 6 h before use. The ABTS⁺ solution was diluted with ethanol to an absorbance of 0.7 (± 0.02) at 734 nm. Then, 1 mL of the working solution was mixed with 10 μ L of EO for 30 s and the absorbance was read at 734 nm after another 20 s using a Jenway 6310 spectrophotometer (Keison, Chelmsford, Essex, UK). An ABTS⁺ solution without EO was used as a control. The results were expressed as μ M eq Trolox mL⁻¹ EO. The assay was performed in triplicate.

2.6 Phytotoxicity Test

A dose-response assay was carried out following Vitalini et al. [15], with some modifications. 2 or 20 μ L of EO was placed in an aluminum container inside 9 cm diameter Petri dishes to avoid direct contact with the seeds distributed on a layer of filter paper wetted with 4 ml of deionized water. The Petri dishes were

prepared in a vertical laminar flow hood and sealed with a double layer of parafilm to prevent the volatile compound escape. Then, they were transferred in a growth chamber at 16/8 h of light/dark photoperiod with a temperature of 25/18 °C for 7 days. The experimental design was as follows: 15 seeds of *L. multiflorum* or *S. alba* × 2 doses of EO or distilled water as a control × 3 replicates × 2 runs.

The seed germination (radicle ≥ 1 mm) was recorded daily. Root and shoot lengths of *L. multiflorum* and *S. alba* seedlings were measured at the end of the tests, after a week. The collected data were used to calculate four indices.

- (i) Germination percentage (G) = Germinated seed number)/(Seed total number) × 100
- (ii) Coefficient of Velocity of Germination (CVG) = $N_1 + N_2 + \dots + N_i/100 \times N_1 T_1 + \dots + N_i T_i$, where N is the number of seeds germinated every day; T is the number of days from seeding corresponding to N [16]
- (iii) Mean Germination Time (MGT) = $(\sum D \times \text{Germinated seed number})/(\sum \text{Germinated seed number})$, where D is the number of days from the beginning of germination, plus the number of seeds germinated on day D [17]
- (iv) Seedling Vigor Index (SVI) = (Mean Root length + Mean Shoot length) × Germination % [18].

3. Results and Discussion

3.1 Chemical analysis

The GC-MS technique was used to describe the chemical profile of *P. ostruthium* leaf EO. In total, 32 compounds, accounting for 99.6% of the total composition, were identified, and their average percentage values are shown in Table 1. EO was characterized by a sesquiterpene content exceeding the monoterpenic one and the main two components were caryophyllene oxide (20.7%) and spathulenol (17.2%), followed by cubenol (8.7%), δ-cadinene (6.1%), humulene epoxide II (5.6%) and *cis*-*p*-mentha-2,8-dien-1-ol (5.4%). All detected monoterpenes ranged from 0.1 to 0.9%. Two phthalates, namely diisobutyl phthalate (4.6%) and diethyl phthalate (1.3%), two fatty acids such as decanoic acid (0.3%) and palmitic acid (0.7%), and a diterpenoid, 13-

epimanool (3.8%), were also found. The GC-FID chromatogram was reported in Fig. 1.

Table 1. Chemical composition (percentage values ± standard deviation) of *P. ostruthium* leaf essential oil.

| Compounds ¹ | LRI ² | LRI ³ | Area (%) |
|---|------------------|------------------|-------------|
| α-Pinene | 938 | 945 | 0.4±0.03 |
| cis-Linalool oxide | 1052 | * | 0.1±0.02 |
| Linalool | 1092 | 1088 | 0.4±0.02 |
| Chrysanthrone | 1106 | 1103 | 1.0±0.03 |
| <i>Cis</i> - <i>p</i> -mentha-2,8-dien-1-ol | 1120 | 1116 | 5.4±0.05 |
| Dehydrolinalool | 1121 | 1116 | 0.5±0.02 |
| Trans- <i>p</i> -mentha-1(7),8-dien-2-ol | 1171 | 1165 | 0.9±0.02 |
| α-Terpineol | 1183 | 1182 | 0.7±0.04 |
| <i>p</i> -Cymen-8-ol | 1186 | 1185 | 0.7±0.03 |
| Trans-sabinene hydrate | 1190 | * | 1.0±0.05 |
| 2,5-Bornanedione | 1268 | 1264 | 0.5±0.03 |
| Decanoic acid | 1355 | 1353 | 0.3±0.02 |
| α-Copaene | 1378 | 1388 | 0.5±0.03 |
| (-)-β-Bourbonene | 1392 | 1390 | 0.5±0.02 |
| Trans-geranyl acetone | 1430 | 1432 | 0.2±0.02 |
| β-Caryophyllene | 1431 | 1434 | 3.5±0.02 |
| <i>Cis</i> -muurola-3,5-diene | 1451 | 1447 | 0.8±0.02 |
| Humulene | 1475 | 1473 | 3.0±0.03 |
| Germacrene D | 1493 | 1489 | 0.6±0.03 |
| Valencene | 1517 | 1515 | 2.7±0.02 |
| δ-Cadinene | 1521 | * | 6.1±0.02 |
| <i>epi</i> -Cubebol | 1530 | * | 3.4±0.03 |
| Diethyl phthalate | 1555 | 1551 | 1.3±0.02 |
| Spathulenol | 1578 | 1571 | 17.2±0.05 |
| Caryophyllene oxide | 1583 | 1580 | 20.7±0.04 |
| Humulene epoxide II | 1615 | * | 5.6±0.02 |
| Cubenol | 1635 | 1631 | 8.7±0.02 |
| Ledene oxide II | 1678 | * | 3.1±0.04 |
| Hexahydrofarnesyl acetone | 1855 | 1846 | 0.7±0.06 |
| Diisobutyl phthalate | 1868 | 1871 | 4.6±0.03 |
| Palmitic acid | 1978 | 1973 | 0.7±0.03 |
| 13-Epimanool | 2060 | 2056 | 3.8±0.03 |
| Total | | | 99.6 |
| Monoterpenoids | | | 6.4 |
| Sesquiterpenoids | | | 77.1 |
| Diterpenoids | | | 3.8 |
| Others | | | 12.3 |

¹The components are reported according to their elution order on apolar column; ²Linear Retention Indices measured on apolar column; ³Linear Retention Indices from literature; * LRI not available; ⁴Percentage mean values of *P. ostruthium* EO components.




Figure 1. GC-FID Chromatogram of *P. ostruthium* leaf essential oil.

The prevalence of sesquiterpenoid compounds in *P. ostruthium* leaf EO composition confirmed the data previously published in the only work present in the literature where Cisowski and co-authors [19] analyzed, in addition to the rhizome EO, the EO obtained from the herb (it is not better specified, the whole aerial parts including the leaves are assumed) of *P. ostruthium* collected in the Sudety mountains (Karpacz area, Poland). In that case, the main components were β -caryophyllene (16.1%), α -humulene (15.8%) and germacrene D (9.6%), and the monoterpenoid fraction was more abundant than that of the *P. ostruthium* EO under investigation (25.9% vs 6.4%). Nevertheless, many compounds, both mono- and sesquiterpenoids, were found to be common, albeit, in some cases, in significantly different amounts. Differently, caryophyllene oxide was identified as the main compound (23.1%) in the EO of *P. austriacum* leaves, followed by germacrene D and (E)-caryophyllene [20]. Leaf EOs obtained from other

Peucedanum species such as *P. zenkeri* Engl., *P. petiolare* Boiss., *P. officinale* L. and *P. cervaria* (L.) Lapeyr. were instead characterized by a higher content of monoterpenes, mainly sabinene, α - and β -pinene, limonene, myrcene and β -phellandrene [21-24].

3.2 *P. ostruthium* radical scavenging activity

The values reported in Table 2 showed a greater ability of the *P. ostruthium* leaf EO to scavenge the ABTS⁺ radical cation rather than the DPPH[·] free radical. In terms of percentage inhibition, ABTS⁺ was reduced by 2.2 times more than DPPH[·] (91.0% vs 41.2%). The difference could be attributed to the different solubility of EO in the test systems and can be considered as an indication of the relative reactivity of its active compounds as it occurs for resinous exudates or other extracts [25, 26].

Few previous reports have documented the antiradical and/or antioxidant activity of *Peucedanum* species. Masuda et al. [27] reported a weak ability of

Table 2. Antiradical activity of essential oil from *P. ostruthium*

| | ABTS (μ M Trolox eq/mL) | DPPH (μ M Trolox eq/mL) |
|----|---------------------------------|---------------------------------|
| EO | 2.02 \pm 0.00 | 0.77 \pm 0.01 |

Values are mean \pm standard deviation.

the methanolic extract of *P. japonicum* leaves to inhibit DPPH. A few years later, Hisamoto and collaborators [28] obtained a very active butanol fraction from the leaves of the same species. Hence, some compounds including rutin and three isomers of caffeoylquinic acid were isolated and found to be the major antioxidant constituents. Only the antiradical activity of the *P. litorale* EO seems to have been studied [29]. The data shows that a comparison is rather difficult. The type of sample, the different origin, the method of extraction and analysis are just some factors that influence its bioactivity.

3.3 *P. ostruthium* phytotoxicity

Appreciable results were obtained from screening of the phytotoxic activity of *P. ostruthium* leaf EO against both target species at the two tested doses (Table 3). It was found to be less effective in influencing seed germination (reduced by 39.6% and 33.3%, respectively) than seedling growth. In this case, the 2 μ L dose had greater effects towards *S. alba* (SVI, -36.8% vs -10.9%; root length, -31.8% vs -8.3%; shoot length, -24.1% vs -7.7%) while the 20 μ L dose showed a better effect against *L. multiflorum* (SVI, -90.7% vs -76.6%; root length, -90.4% vs -67.6%; shoot length, -76.4% vs -60.4%). Under the effect of 20 μ L of EO, the CVG and MGT indices of *L. multiflorum* were also the most affected, decreasing by 64.2% and increasing by 16.7%, respectively.

These results supported previously published data on

the phytotoxic potential of *P. ostruthium* [30] and proved its herbicidal action against dicotyledonous species. In fact, the aqueous extracts and powders of inflorescences, leaves and rhizomes of *P. ostruthium* from the same geographical area were able to significantly reduce both the germination and the development of two important monocotyledonous weeds, namely *Echinochloa oryzoides* (Ard.) Fritsch and *L. multiflorum*. 5-Caffeoylquinic acid was the main phenolic component in all plant organs, but other chemical classes such as chlorogenic acids, flavonol glycosides, coumarins, and furanocoumarin glycosides were detected. Among the compounds identified in *P. ostruthium* leaf EO, some in particular might be responsible for the inhibitory activity. Caryophyllene oxide and spathulenol, the two most abundant, have been reported to possess a variety of biological properties [31,32], including allelopathic one [33-35]. Besides them, cubenol was responsible for the allelopathic effect of *Sinapis arvensis* var. *orientalis* EO [36], δ -cadinene was identified as one of the exudates released by the roots of *Chrysanthemoides monilifera* spp. *rotundata* (DC.) T. Norl and found in the extracts of its growth soil characterized by phytotoxic activity [37].

Other sesquiterpenoids of the EO were recognized as allelochemicals (e.g., β -caryophyllene) or detected as the major compounds of EOs with significant allelopathic behavior (e.g., humulene) [38,39]. Allelopathy of monoterpenes is also documented [40]. For example, α -pinene is a compound present in EOs with high activity [41]. This component has an important ecological role on the plant-plant allelopathic interference. Pinenes were found to reduce chlorophyll content in leaves, cell respiration,

Table 3. Germination and growth parameters of two target species *L. multiflorum* and *S. alba* under the phytotoxic effects of *P. ostruthium* essential oil.

| Target species | EO (μ L) | G (%) | CVG | MGT | SVI | Root (mm) | Shoot (mm) |
|-----------------------|---------------|-----------------|----------------|---------------|------------------|-----------------|-----------------|
| <i>L. multiflorum</i> | 0 | 96.0 \pm 4.0 | 109 \pm 7.0 | 4.8 \pm 0.0 | 12538 \pm 1183 | 75.9 \pm 13.7 | 54.7 \pm 11.2 |
| | 2 | 93.0 \pm 0.0 | 93 \pm 5.5 | 5.0 \pm 0.0 | 11169 \pm 946 | 69.6 \pm 8.5 | 50.5 \pm 1.2 |
| | 20 | 58.0 \pm 14.0 | 39 \pm 3.0 | 5.6 \pm 0.1 | 1172 \pm 244 | 7.3 \pm 1.9 | 12.9 \pm 1.9 |
| <i>S. alba</i> | 0 | 90.0 \pm 5.0 | 122 \pm 6.2 | 4.1 \pm 0.1 | 4999 \pm 384 | 34.3 \pm 6.7 | 21.2 \pm 4.7 |
| | 2 | 80.0 \pm 5.0 | 112 \pm 10.0 | 4.2 \pm 0.1 | 3160 \pm 195 | 23.4 \pm 4.6 | 16.1 \pm 5.0 |
| | 20 | 60.0 \pm 0.0 | 68 \pm 3.00 | 4.4 \pm 0.1 | 1170 \pm 112 | 11.1 \pm 1.6 | 8.4 \pm 1.0 |

Values are mean \pm standard deviation. G%, Germination percentage; CVG, Coefficient of Velocity of Germination; MGT, Mean Germination Time, SVI, Seedling Vigor Index

enzymatic activity of proteases, α - and β -amylases as well as root and coleoptile length [42]. Therefore, the phytotoxicity of *P. ostruthium* leaf EO could be linked to the presence of both monoterpenes and sesquiterpene, whose effectiveness can in some cases be improved by a synergistic interaction [41].

4. Conclusions

To conclude, this is one of the very few works on *P. ostruthium* and the first on the antiradical and phytotoxic activity of its EO. The obtained results are promising and deserve to be corroborated with further tests. *P. ostruthium* is confirmed as a good source of compounds, which have proved to be useful both as antioxidants and as allelochemicals.

Author Contributions

Conceptualization, S.V. and M.I.; Methodology, S.V., S.G. and M.I.; Validation, S.V. and S.G.; Investigation, S.V. and S.G.; Resources, S.G. and M.I.; Data Curation, S.V. and S.G.; Writing – Original Draft Preparation, S.V. and S.G.; Writing – Review & Editing, S.V., S.G. and M.I.

Acknowledgements and Funding

This research received no specific grant from any funding agency.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Tutin T.G.; Heywood V.H.; Burges N.A.; Moore D.M.; Valentine D.H.; Walters S.M.; Webb D.A. *Flora Europaea*, 2nd ed. Cambridge University Press, Cambridge, UK, **1993**.
2. Pignatti, S. *Flora d'Italia*, 2nd ed. Edagricole, Bologna, Italy, **2017**.
3. Vogl S.; Picker, P.; Mihaly-Bison J.; Fakhrudin N.; Atanasov, A.G.; Heiss E.H.; Wawrosch C.; Reznicek, G.; Dirsch V.M.; Saukel J.; Kopp B. Ethnopharmacological *in vitro* studies on Austria's folk medicine-An unexplored lore *in vitro* anti-inflammatory activities of 71 Austrian traditional herbal drugs. *J. Ethnopharmacol.* **2013**, *149*, 750–771.
4. Vitalini S.; Puricelli C.; Mikerezi I.; Iriti M. Plants, people and traditions: ethnobotanical survey in the Lombard Stelvio National Park and neighbouring areas (Central Alps, Italy). *J. Ethnopharmacol.* **2015**, *173*, 435–458.
5. Danna C.; Poggio L.; Smeriglio A.; Mariotti M.; Cornara L. Ethnomedicinal and ethnobotanical survey in the Aosta Valley side of the Gran Paradiso National Park (Western Alps, Italy). *Plants* **2022**, *11*, 170.
6. Sarkhail P. Traditional uses, phytochemistry and pharmacological properties of the genus *Peucedanum*: a review. *J. Ethnopharmacol.* **2014**, *156*, 235–270.
7. Alavi S.H.R.; Yassa N.; Fazeli, M.R. Chemical constituents and antibacterial activity of essential oil of *Peucedanum ruthenicum* M. Bieb. fruits. *Iran. J. Pharm. Sci.* **2005**, *1*, 217–222.
8. Yang E.-J.; Kim S.-S.; Oh T.-H.; Song G.; Kim K.-N.; Kim J.-Y.; Lee N.H.; Hyun C.-G. *Peucedanum japonicum* and *Citrus unshiu* essential oils inhibit the growth of antibiotic-resistant skin pathogens. *Ann. Microbiol.* **2009**, *59*, 623–628.
9. Lim H.; Shin S. Study on the essential oils from the roots of *Angelica decursiva* and *Peucedanum praeruptorum*. *Kor. J. Pharmacogn.* **2012**, *43*, 291–296.
10. Khruengsai S.; Sripahco T.; Rujanapun N.; Charoensup R.; Pripdeevech P. Chemical composition and biological activity of *Peucedanum dhana* A. Ham essential oil. *Sci. Rep.* **2021**, *11*, 19079.
11. Sun J.; Feng Y.; Wang Y.; Li J.; Zou K.; Liu H.; Liu H.; Hu Y.; Xue Y.; Yang L.; Du S.; Wu Y. Investigation of pesticidal effects of *Peucedanum terebinthinaceum* essential oil on three stored-product insects. *Rec. Nat. Prod.* **2020**, *14*, 177–189.
12. Vitalini S.; Iriti M.; Vaglia V.; Garzoli S. Chemical investigation and dose-response phytotoxic effect of essential oils from two gymnosperm species (*Juniperus communis* var. *saxatilis* Pall. and *Larix decidua* Mill.). *Plants* **2022**, *11*, 1510.
13. Vitalini, S.; Iriti, M.; Garzoli, S. GC-MS and SPME-GC/MS Analysis and bioactive potential evaluation of essential oils from two *Viola* species belonging to the *V. calcarata* complex. *Separations* **2022**, *9*, 39.
14. Iriti M.; Vitalini S.; Apostolides N.A.; El Beyrouthy M. Chemical composition and antiradical capacity of essential oils from Lebanese medicinal plants. *J. Ess. Oil Res.* **2014**, *26*, 466–472.

15. Vitalini S.; Orlando F.; Iriti M. Selective phytotoxic activity of eugenol towards monocot and dicot target species. *Nat. Prod. Res.* **2021**, *36*, 1659–1662.

16. Al-Mudaris, M. Notes on various parameters recording the speed of seed germination. *Der Trop.* **1998**, *99*, 147–154.

17. Ellis R.A., Roberts E.H. The quantification of ageing and survival in orthodox seeds. *Seed Sci. Technol.*, **1981**, *9*, 373–409.

18. Abdul-Baki A.A., Anderson J.D. Vigour determination in soybean seed by multiple criteria. *Crop Sci.* **1973**, *1*, 630–633.

19. Cisowski W.; Sawicka U.; Mardarowicz M.; Asztemborska M.; Łuczkiewicz M. Essential oil from herb and rhizome of *Peucedanum ostruthium* (L. Koch.) ex DC. *Z. Naturforsch. C*, **2001**, *56*, 930–932.

20. Jovanović O.P.; Zlatković B.K.; Simonović S.R.; Đorđević A.S.; Palić I.R.; Stojanović G.S. (2013). Chemical composition and antibacterial activity of the essential oils isolated from leaves and fruits of *Peucedanum austriacum* (Jacq.) WDJ Koch. *J. Ess. Oil Res.* **2013**, *25*, 129–137.

21. Menut C.; Mve-Mba C.E.; Lamaty G.; Zollo P.H.A.; Tchoumbougnang F.; Bessiere J.M. Aromatic plants of tropical Central Africa. XVIII. Essential oils of leaf and rhizome of *Peucedanum zenkeri* Engl. from Cameroon. *J. Ess. Oil Res.* **1995**, *7*, 77–79.

22. Mirza M.; Najafpour Navaei M.; Dini M. Chemical composition of the essential oils from the rhizome, leaf and seed of *Peucedanum petiolare* (DC.) Boiss. *Flavour Frag. J.* **2005**, *20*, 196–198.

23. Figuérèdo G.; Chalchat J.C.; Petrovic S.; Maksimovic Z.; Gorunovic M.; Boza P.; Radic J. Composition of essential oils of flowers, leaves, stems and rhizome of *Peucedanum officinale* L. (Apiaceae). *J. Ess. Oil Res.* **2009**, *21*, 123–126.

24. Chizzola R. Composition of the essential oils from *Peucedanum cervaria* and *P. alsaticum* growing wild in the urban area of Vienna (Austria). *Nat. Prod. Commun.* **2012**, *7*, 1934578X1200701126.

25. Lissi E.A.; Modak B.; Torres R.; Escobar J.; Urzua A. Total antioxidant potential of resinous exudates from *Heliotropium* species, and a comparison of the ABTS and DPPH methods. *Free Rad. Res.* **2009**, *30*, 471–477.

26. Xie Q.; Liu Z.; Li Z. Chemical composition and antioxidant activity of essential oil of six *Pinus* taxa native to China. *Molecules* **2015**, *20*, 9380–9392.

27. Masuda T.; Yonemori S.; Oyama Y.; Takeda Y.; Tanaka T.; Andoh T.; Shinohara A.; Nakata M. Evaluation of the antioxidant activity of environmental plants: Activity of the leaf extracts from seashore plants. *J. Agric. Food Chem.* **1999**, *47*, 1749–1754.

28. Hisamoto M.; Kikuzaki H.; Ohigashi H.; Nakatani N. Antioxidant compounds from the leaves of *Peucedanum japonicum* Thunb. *J. Agric. Food Chem.* **2003**, *51*, 18, 5255–5261.

29. Iskakova Z.B.; Suleimen Y.M.; Dudkin, R.V.; Gorovoy, P.G. Constituent of cytotoxic and antiradical activity of essential oil *Peucedanum litorale*. In 12th International Symposium on the Chemistry of Natural Compounds **2017**, pp. 107–107.

30. Vitalini S.; Palmioli A.; Orlando F.; Scari G.; Auiroldi C.; De Noni I.; Bocchi S.; Iriti M. Phytotoxicity, nematicidal activity and chemical constituents of *Peucedanum ostruthium* (L.) W.D.J.Koch (Apiaceae). *Ind. Crops Prod.* **2021**, *166*, 113499.

31. do Nascimento K.F.; Figueira Moreira F.M.; Santos J.A.; Leite Kassuya C.A.; Rosa Ernesto de Carvalho J.; Nazari Formagio A.S. Antioxidant, antiinflammatory, antiproliferative and antimycobacterial activities of the essential oil of *Psidium guineense* Sw. and spathulenol. *J. Ethnopharmacol.* **2017**, *210*, 351–358.

32. Gyrdymova Y.V.; Rubtsova S.A. Caryophyllene and caryophyllene oxide: a variety of chemical transformations and biological activities. *Chem. Pap.* **2022**, *76*, 1–39.

33. Quintana N.; El Kassis E.G.; Stermitz F.R.; Vivanco J.M. Phytotoxic compounds from roots of *Centaurea diffusa* Lam. *Plant Signal. Behav.* **2009**, *4*, 9–14.

34. Verdeguer M., M.A. Blazquez and Boira H. Phytotoxic effects of *Lantana camara*, *Eucalyptus camaldulensis* and *Eriophyllum africanus* essential oils in weeds of Mediterranean summer crops. *Biochem. Syst. Ecol.* **2009**, *37*, 362–369.

35. Razavi S. M. Chemical and allelopathic analyses of essential oils of *Prangos pabularia* Lindl. from Iran. *Nat. Prod. Res.* **2012**, *26*, 2148–2151.

36. Sharifi-Rad J.; Miri A.; Sharifi-Rad M., Sharifi-Rad R.; Sharifi-Rad M. Allelopathic effects of essential oils from *Sinapis arvensis* L. aerial part on germination and seedling growth of medicinal plants and weeds. *Int. J. Biosci.* **2014**, *5*, 135–140.

37. Ens E.J.; Bremner J.B.; French K.; Korth J. Identification of volatile compounds released by roots of an invasive plant, bitou bush (*Chrysanthemoides monilifera* spp. *rotundata*), and their inhibition of native seedling growth. *Biol. Invasions* **2009**, *11*, 275–287.

38. Mota M.S.C.S.; Silva R.S.; Silva G.A.; Picanco M.C.; Mesquita A.L.M.; Pereira R.C.A. Potential of allelochemicals from basil (*Ocimum micranthum* Willd.) to control whitefly (*Aleurodicus cocois* (Curtis 1846)) in cashew nut crop (*Anacardium occidentale* L.). *Allelopathy J.* **2017**, *40*, 197–208.

39. Abd-ElGawad A.M.; Bonanomi G.; Al-Rashed S.A.; Elshamy A.I. *Persicaria lapathifolia* essential oil: Chemical constituents, antioxidant activity, and allelopathic effect on the weed *Echinochloa colona*. *Plants* **2021**, *10*, 1798.

40. Macías F.A.; Mejías F.J.; Molinillo J.M. Recent advances in allelopathy for weed control: from knowledge to applications. *Pest Manag. Sci.* **2019**, *75*, 2413–2436.

41. Amri I.; Hamrouni L.; Hanana M.; Jamoussi B. Reviews on phytotoxic effects of essential oils and their individual components: news approach for weed management. *Int. J. Appl. Biol. Pharm. Technol.* **2013**, *4*, 96–114.

42. Chowhan N.; Singh H.P.; Batish D.R.; Kohli R.K. Phytotoxic effects of α -pinene on early growth and associated biochemical changes in rice. *Acta Phys. Plant.* **2011**, *33*, 2369–2376.